

# Chapter 9

## Sequences and Series

### § 9.1 Infinite Sequences

An infinite sequence (or simply a sequence) is simply an infinite (ordered) list of numbers

$$s_1, s_2, s_3, \dots, s_n, \dots$$

The individual numbers  $s_n$  are called the terms of the sequence ( $s_1$  is the first term,  $s_2$  the second term and so on).

A common way of writing a sequence is

$$\{s_n\}_{n=1}^{\infty}$$

(not used in the textbook, but very standard nonetheless).

## Examples

$$1) S_n = 1, \text{ for every } n \geq 1$$

$$\text{Get } \{1\}_{n=1}^{\infty} = 1, 1, 1, 1, \dots$$

$$2) S_n = (-1)^n, \quad n \geq 1$$

$$\text{Get } \{(-1)^n\}_{n=1}^{\infty} = -1, 1, -1, 1, -1, \dots$$

$$3) S_n = \frac{1}{n}, \quad n \geq 1$$

$$\text{Get } \left\{\frac{1}{n}\right\}_{n=1}^{\infty} = 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$$

The above are all examples of one of the main ways of representing a sequence - using a formula in  $n$ .

A common problem is to find a suitable formula when you're given the first several terms of the sequence.

Ex

a) 3, 6, 12, 24, 48, 96, 192, 384, ...

What we notice here first is that each term is double the previous one. This suggests that  $2^n$  should be part of our formula.

Also, the first term is 3 while all the terms are divisible by 3. This suggests there should be a 3 in our formula.

Try 
$$S_n = 3(2)^n$$

This fits all the terms we were given and so (as far as we can tell) this is a formula for our sequence.

$$b) \quad \frac{7}{2}, \frac{7}{5}, \frac{7}{8}, \frac{7}{11}, \frac{1}{2}, \frac{7}{17}, \dots$$

Here the  $\frac{1}{2}$  term looks like the rest  
but if we write  $\frac{1}{2}$  as  $\frac{7}{14}$ , we get

$$\frac{7}{2}, \frac{7}{5}, \frac{7}{8}, \frac{7}{11}, \frac{7}{14}, \frac{7}{17}, \dots$$

and the pattern is more evident.

The general term should obviously be  
a fraction whose numerator is 7.

Also the denominator clearly increases  
by 3 as we go from one term to the  
next while the first term is 2.

Suggests the denominator is of the form

$$3n + k$$

and since at 1, we must have  $3(1) + k = 2$ ,  
we get  $k = -1$ , and so we get  $3n - 1$ .

A formula which fits our data is then

$$S_n = \frac{7}{3n-1}, \quad n \geq 1.$$

Consider now the sequence

1, 1, 2, 3, 5, 8, 13, 21, 34, 55, ...

This is the famous Fibonacci Sequence where each term is the sum of the preceding two. More precisely,

$$s_1 = 1, s_2 = 1, s_n = s_{n-1} + s_{n-2}, n \geq 3.$$

This is an example of the second main way of defining a sequence which is recursively where the formula for each term (except possibly the first few) involves previous terms. We generally also need to specify the first few terms to get the recursion started (eg. for Fibonacci, we needed to first give  $s_1$  and  $s_2$ ).

Ex. Find a recursive formula for each of the following.

a) 1, 2, 6, 24, 120, 720, 5040, ...

Here we notice that each term  $s_n$  is  $n$  times the previous one. - e.g.

$6 = 3(2)$ ,  $720 = 6(120)$ . A recursive formula is then

$$s_1 = 1, \quad s_n = n s_{n-1}, \quad n > 1.$$

Note that here  $s_n$  is the famous factorial of  $n$ ,  $n!$  where

$$n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1)(n),$$

the product of the first  $n$  natural numbers.

b) 1, 3, 7, 15, 31, 63, 127, ...

Here the first term is 1 while each term is obtained from the last one by doubling and adding 1 (e.g.

$$7 = 2(3) + 1, \quad 127 = 2(63) + 1).$$

A recursive defn which fits is then

$$S_1 = 1, \quad S_n = 2S_{n-1} + 1, \quad n > 1.$$

# Convergence of Sequences

Consider the sequence

$$\left\{ \frac{1}{n} \right\}_{n=1}^{\infty} = 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$$

The bigger  $n$  is, the smaller  $\frac{1}{n}$  becomes and we can clearly make  $\frac{1}{n}$  as small as we like by taking  $n$  large enough.

This motivates the following defn.

Defn. The sequence  $\{s_n\}_{n=1}^{\infty} = s_1, s_2, s_3, \dots$  has a limit  $L$ , written

$$\lim_{n \rightarrow \infty} s_n = L$$

if  $s_n$  is as close to  $L$  as we please whenever  $n$  is sufficiently large. If a limit  $L$  exists, we say the sequence converges to  $L$ . Otherwise, if no limit exists, we say the sequence diverges.

# Important Examples

$$I \quad S_n = x^n$$

If  $|x| < 1$ , then  $\{x^n\}_{n=1}^{\infty}$  converges to 0

e.g. if  $x = \frac{1}{2}$ , we get  $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$

If  $|x| > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  diverges

e.g. if  $x = -3$ , we get  $-3, 9, -27, 81, \dots$

If  $x = 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is

just the sequence  $1, 1, 1, 1, \dots$

which (trivially) converges to 1.

Finally, if  $x = -1$ , then  $\{x_n\}_{n=1}^{\infty}$  is

the sequence  $-1, 1, -1, 1, -1, 1, \dots$

which diverges.

$$\text{II} \quad S_n = \frac{1}{n^p} \quad (p \text{ fixed}).$$

If  $p > 0$ , then  $\left\{ \frac{1}{n^p} \right\}_{n=1}^{\infty}$  clearly converges to 0 (e.g. if  $p = \frac{1}{2}$ , we get

$$1, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{3}}, \frac{1}{2}, \frac{1}{\sqrt{5}}, \dots$$

If  $p < 0$  then  $\frac{1}{n^p} = n^{-p}$  gets

large as  $n \rightarrow \infty$  since  $-p > 0$ . In this case  $\left\{ \frac{1}{n^p} \right\}_{n=1}^{\infty}$  diverges.

Finally, if  $p = 0$   $\frac{1}{n^0} = 1$  for every  $n$  and so we just have the sequence

$$1, 1, 1, \dots$$

which is convergent to 1.

Ex.

a)  $S_n = (0.8)^n$

This is of the type in Ex I with  $|0.8| < 1$  and so converges to 0.

b)  $S_n = \frac{1 - e^{-n}}{1 + e^{-n}}$

Since  $|e^{-1}| < 1$ ,  $\lim_{n \rightarrow \infty} e^{-n} = 0$  and so

$$\lim_{n \rightarrow \infty} S_n = \frac{1 - 0}{1 + 0} = 1$$

c)  $S_n = \frac{1 + e^n}{1 - e^n}$

Here we divide above and below by  $e^n$  and remember that  $\frac{1}{e^n} = e^{-n}$  to get

$$S_n = \frac{e^{-n} + 1}{e^{-n} - 1}$$

Similarly to b), we then see that  $\lim_{n \rightarrow \infty} S_n = -1$ .

## Facts about Sequences

A sequence  $\{s_n\}_{n=1}^{\infty}$  is called bounded if we can find numbers  $K$  and  $M$  such that

$$K \leq s_n \leq M \quad \text{for every } n.$$

### Fact

A convergent sequence is bounded.

Idea. If  $\{s_n\}$  is convergent with limit  $L$ , then eventually  $s_n$  is close to  $L$  (e.g. between  $L-1$  and  $L+1$ ) for all sufficiently large  $n$ . and so at least these later terms of the sequence are bounded.

The remaining terms are the early ones, but as there are only finitely many of them, they will not affect whether the sequence is bounded.

A sequence  $\{s_n\}_{n=1}^{\infty}$  is called monotone if it is either increasing, i.e.

$$s_n < s_{n+1} \quad \forall n \geq 1$$

or it is decreasing

$$s_n > s_{n+1} \quad \forall n \geq 1.$$

Fact.

If  $\{s_n\}_{n=1}^{\infty}$  is a sequence which is bounded and monotone, then it is convergent.

Ex. If  $s_n = (1 + \frac{1}{n})^n$ , one can show that

$s_n$  is increasing and that  $s_n < 3$  for every  $n$ , so that we have a bounded sequence.

By the above fact  $s_n$  is then convergent.

In fact we already know the limit of this sequence is  $e$  using l'Hôpital's rule.